# **POSTOPERATIVE INFORMATION**

# JEFFREY WONG, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE SCOS ORTHOPEDIC SPECIALISTS 714.500.5056 OFFICE 949.900.2116 FAX CA License: A114599

PATIENT STICKER

# POST-OPERATIVE INSTRUCTIONS -

# DEBRIDEMENT AND REPAIR OF COMMON EXTENSOR OR FLEXOR (TREATMENT OF EPICONDYLITIS):

#### WOUND CARE

- □ You may remove the Operative Dressing on Post-Op Day #3.
- Apply sterile gauze to the wound. Change the gauze daily. Please <u>do not use</u> Bacitracin or other ointments under the bandage. There are white tapes directly on the skin, leave these on until seen by Dr. Wong. Loosely re-wrap the ace bandage for comfort.
- □ Use Ice 3-4 times per day for the first 3-4 days, then as needed for pain relief.
- □ A sling will be placed on your arm in the recovery room. You may remove it when at home and straighten your arm, but you can wear it in public places for about 7 days.
- □ Rest with a pillow under your elbow.
- □ You may shower on Post-Op Day #3. Please cover the wound with plastic wrap and secure it to your skin with tape. You may remove the sling for showering. Gently pat the area dry. <u>Do not soak</u> the elbow in water. <u>Do not go swimming</u> in the pool or ocean until clear by Dr. Wong.

#### **EXERCISES**

- □ Avoid lifting with your operated arm until seen in the office for wound check and brace application.
  - <u>Gentle</u> Elbow Flexion / Extension (gentle bending and straightening)
  - Hand / Wrist Range of Motion Exercises and Grip Strengthening

#### POST-OP

- $\hfill\square$  You will be given two prescriptions for pain relief for you to use post-operatively:
  - NAPROXEN, an anti-inflammatory medication, is to be used twice a day with breakfast and dinner, for the first ten days after surgery.
    - \*\* IF YOU HAVE AN ULCER HISTORY OR DIFFICULTY USING ANTI-INFLAMMATORIES OR ASPIRIN >> DO NOT USE
  - NORCO or PERCOCET, a strong narcotic, to be used only on an <u>"as needed"</u> basis for pain in addition to the Naprosyn.
- □ If you have any adverse effects with the medications, please call our office.
- □ If you develop a Fever (101.5), Redness or Drainage from the surgical incision site, please call our office to arrange for an evaluation.

#### FOLLOW-UP

□ Please call the office to schedule a follow-up appointment for wound check, 10-14 days postoperatively.

### IF YOU HAVE ANY QUESTIONS, PLEASE FEEL FREE TO CALL OUR OFFICE.