

# POSTOPERATIVE INFORMATION

## JEFFREY WONG, MD

ORTHOPAEDIC SURGERY AND SPORTS MEDICINE  
SCOS ORTHOPEDIC SPECIALISTS  
714.500.5056 OFFICE  
949.900.2116 FAX  
CA License: A114599

PATIENT STICKER

## POST-OPERATIVE INSTRUCTIONS – MENISCAL REPAIR

### DAY OF SURGERY:

1. A Cryocuff (cold therapy device) will be placed on your knee in the recovery room. For the first 24 hours following surgery, keep the cuff on as much as possible except when you go to bed. This will help reduce the swelling and minimize the pain. However, you will need to continuously change the water so that it remains cold.
2. Your brace will be applied in the operating room. You must wear the brace locked straight for 6 weeks when walking and sleeping. You may unlock it for short periods when you are sitting 2 days after the surgery.
3. There are exercises included that you can begin on the first or second postoperative day. To protect the meniscus do not bend the knee past 90 degrees for the first six weeks after your surgery.
4. Keep your leg elevated to decrease swelling, which will then in turn decrease your pain. When in bed, your leg should be straight with a pillow under the calf, NOT under the knee.
5. Prescriptions for pain medication will be given to you when you leave. You should avoid taking pain medications on an empty stomach, as it will make you nauseous. Otherwise, use the pain medication as needed. Norco as needed for pain. Colace as needed for constipation. Zofran as needed for nausea. Aspirin daily for 2 weeks to prevent blood clots.

### DAY 1 AND UNTIL FIRST POST-OP VISIT:

1. The cryocuff will aid in decreasing pain and swelling. Apply the cryo-cuff for 20-30 minutes at least 4 times per day for the first few days. Then use it 2-3 times a day. You may use the cryo-cuff more frequently if you are having continued pain and swelling.
2. Remove the bandages 72 hours after surgery and apply gauze to the wounds. Do not remove the white steri-strip tape over the incisions. Re-apply clean gauze over the incisions until the wounds are completely dry. Do not put anything including bacitracin or ointments on or near the incisions. Use the ace wrap to decrease swelling but do not wrap it too tightly.
3. You may shower 72 hours after surgery but **Do Not** get the sutures wet. Wrap your knee with plastic wrap and secure it with tape around your thigh. Then, place the brace on and cover it using a cast bag

## POSTOPERATIVE INFORMATION

or a garbage bag with a hole in the bottom for your foot. Secure it with tape around your ankle and thigh. If your incisions do get wet, pat them dry.

4. Use the crutches while standing and walking so that you are only putting partial weight on your operated leg for the first 2 weeks. Ok to place as much weight as you can tolerate.
5. Continue doing the home exercises 2x per day. Begin physical therapy 2 weeks after your surgery.
6. Do not drink alcoholic beverages or take illicit drugs when taking pain medications.
7. You may return to sedentary work/school in the next couple of days when you feel up to it. You will need to keep your leg elevated as much as possible.
8. Call the office tomorrow with any questions, and make an appointment to have your incisions checked in 14 days.

## MISCELLANEOUS INFORMATION

1. There may be some bleeding and fluid leaking from the incision site. This is normal after this type of surgery. This may continue for 24-36 hours. You may change and/or reinforce the bandages as needed. **Do Not** remove the white steri-strip tapes covering the lower incision even if they are slightly wet or bloody.
2. There will be MORE swelling on days 1-3 than there is on the day of surgery. This also is normal. The swelling will decrease with the anti-inflammatory medication, the cryo-cuff and keeping it elevated. The swelling will make it more difficult to bend your knee. As the swelling goes down your motion will become easier.
3. You may develop swelling and bruising that extends from your knee down to your calf and perhaps even to your foot over the next week. Do not be alarmed. This too is normal, and it is due to gravity.
4. There may be some numbness adjacent to the incision site. This may last for 6-12 months.
5. It is also normal to develop a low-grade fever after surgery (up to 100.5°). This can last 2 days after surgery.
6. If there are any questions, please feel free to call the office.