

PHYSICAL THERAPY PRESCRIPTION

Jeffrey Wong, MD

SCOS Orthopedic Specialists
18785 Brookhurst Street, Ste 100
Fountain Valley, CA 92708
(714) 500-5056 Phone
(949) 900-2116 Fax

DATE: _____

PATIENT NAME: _____

D.O.B: _____ MRN: _____

DIAGNOSIS: KNEE PCL TEAR

Right Left Bilateral

SURGERY: KNEE PCL RECONSTRUCTION

DATE OF SURGERY: _____

PHASE I (0-4 WEEKS) – IMMOBILIZATION AND EARLY STRENGTHENING

Goals

- Protect graft and fixation, avoid posterior tibial sag
- Obtain full extension and flexion to 60°

Rehab

- NWB with brace locked in extension
- Full extension allowed with posterior leg supported at all times to prevent tibial sag
- Prone passive flexion < 60°
- **No open chain hamstring strengthening
- Cryotherapy, anti-inflammatory modalities
- Patellar mobilization
- Begin quad strengthening – 3-way leg raises in brace (no flexion), quad sets, electrical stim

PHASE II (5-8 WEEKS) – EARLY STRENGTHENING

Goals

- Maintain full extension and gain flexion to 90° by 6 weeks and 110° by 8 weeks
- Protect graft

Rehab

- Begin partial WB (up to 50%), unlock brace to 30° at 6 weeks and 60° at 8 weeks
- Continue prone passive knee flexion and quadriceps stretching
- Continue patellar mobilization
- Begin SLR with brace locked in extension, progress to mini squats at 6 weeks with brace locked at 30° and at 8 weeks with brace locked at 60°
- Begin single leg balance at 6 weeks in brace
- *No open chain hamstring strengthening*

(continue on next page)

PHYSICAL THERAPY PRESCRIPTION

PHASE III (9-12 WEEKS) – ADVANCED STRENGTHENING

Goals

- Achieve full flexion and extension
- Improved strength, endurance, proprioception

Rehab

- Advance to full WB in brace
- Continue ROM exercises
- Continue strengthening – initiate half squats in brace, step downs, slow progression of multi-plane closed chain exercises but *no open chain hamstring exercises*
- Continue proprioceptive training

PHASE IV (12 WEEKS AND BEYOND) – RETURN TO SPORT

Fit for functional PCL brace

Stationary bike with light resistance

Open chain hamstring strengthening at 4 months

Straight running 6 months

Sport-specific rehab at 9 months

Return to play 9-12 months

Treatment: Eval and Treat or 1 2 3 times per week

Duration: 4 6 weeks

Home Program

**Please send progress notes.

Physician's Signature: _____



NPI #: 1164790507